

Auto

SE/Sem III/CBCS/MECH/M-J-18
AUTO

08/05/2018

Q.P. Code : 39159

(3hours)

[Total marks: 80]

- N.B.** 1) Question No. 1 is compulsory.
2) Answer **any Three** from remaining
3) Figures to the right indicate full marks

1. a) Find Laplace transform of $f(t) = e^{-t} \sin t \cdot \cos 2t$. 5
- b) Show that the set of functions $\cos nx, n = 1, 2, 3 \dots$ is orthogonal on $(0, 2\pi)$. 5
- c) The equations of lines of regression are $x + 2y = 5$ and $2x + 3y = -8$.
Find i) means of x and y , ii) coefficient of correlation between x and y . 5
- d) Evaluate $\int_C (z^2 - 2\bar{z} + 1) dz$ where C is the circle $|z| = 1$. 5
2. a) Using convolution theorem, find the inverse Laplace transform of 6

$$F(s) = \frac{1}{(s^2 + 9)(s^2 + 4)}$$
- b) Obtain Fourier series of $f(x) = |x|$ in $(-\pi, \pi)$ 6
- c) Find the bilinear transformation which maps the points $z = 1, i, -1$ onto the points $w = i, 0, -i$. Hence, find the image of $|z| < 1$ onto the w -plane. 8
3. a) If $v = e^x \sin y$, prove that v is a harmonic function. Also find the corresponding harmonic conjugate function and analytic function. 6
- b) Using Bender-Schmidt method, solve $\frac{\partial^2 u}{\partial x^2} - \frac{\partial u}{\partial t} = 0$, subject to the conditions,
 $u(0, t) = 0, u(5, t) = 0, u(x, 0) = x^2(25 - x^2)$ taking $h = 1$, for 3 minutes. 6
- c) Using Residue theorem, evaluate 8
- i) $\int_0^{2\pi} \frac{d\theta}{2 + \cos\theta}$ ii) $\int_0^\infty \frac{dx}{x^2 + 1}$

[TURN OVER]

4. a) Solve by Crank –Nicholson simplified formula $\frac{\partial^2 u}{\partial x^2} - \frac{\partial u}{\partial t} = 0$,

$u(0, t) = 0, u(1, t) = 2t, u(x, 0) = 0$ taking $h = 0.25$ for two-time steps. 6

b) Obtain the Taylor's and Laurent series which represent the function

$f(z) = \frac{z}{(z-1)(z-2)}$ in the regions, i) $|z| < 1$ ii) $1 < |z| < 2$ 6

c) Solve $(D^2 - 3D + 2)y = 4e^{2t}$ with $y(0) = -3, y'(0) = 5$ where $D \equiv \frac{d}{dt}$ 8

5. a) Find an analytic function $f(z) = u + iv$, 6

$$u = e^{-x}\{(x^2 - y^2) \cos y + 2xy \sin y\}$$

b) Find the Laplace transform of $\frac{\sin at}{t}$. Does the L.T of $\frac{\cos at}{t}$ exist? 6

c) Obtain half range Fourier cosine series of $f(x) = x, 0 < x < 2$. Using Parseval's identity, deduce that – 8

$$\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$$

6. a) Obtain Complex form of Fourier series of $f(x) = e^x, -1 < x < 1$ 6

b) Fit a straight line to the following data, 6

x	100	120	140	160	180	200
y	0.45	0.55	0.60	0.70	0.80	0.85

c) A string is stretched and fastened to two points distance l apart. Motion is started by displacing the string in form $y = a \sin(\pi x / l)$ from which it is released at a time $t = 0$. If the vibrations of a string is given by $\frac{\partial^2 y}{\partial t^2} = c^2 \frac{\partial^2 y}{\partial x^2}$, show that the displacement of a point at a distance x from one end at time t is given by $y(x, t) = a \sin(\pi x / l) \cos(\pi ct / l)$. 8