



- N.B.: 1) Question no.1 is compulsory.  
 2) Attempt any three questions from Q.2 to Q.6.  
 3) Figures to the right indicate full marks.

- Q1. a) Find the Laplace transform of  $e^{-t}t \cosh 2t$ . [5]  
 b) Find the half-range cosine series for  $f(x) = \begin{cases} 1 & , 0 < x < \frac{a}{2} \\ -1 & , \frac{a}{2} < x < a \end{cases}$  [5]  
 c) Find  $\nabla \left( \bar{a} \cdot \nabla \frac{1}{r} \right)$  where  $\bar{a}$  is a constant vector. [5]  
 d) Show that the function  $f(z) = z^3$  is analytic and find  $f'(z)$  in terms of  $z$ . [5]
- Q2. a) Find the inverse Z-transform of  $F(z) = \frac{3z^2 - 18z + 26}{(z-2)(z-3)(z-4)}$ ,  $3 < z < 4$ . [6]  
 b) Find the analytic function whose imaginary part is  $\tan^{-1} \left( \frac{y}{x} \right)$ . [6]  
 c) Obtain Fourier series for the function  $f(x) = \begin{cases} \frac{\pi}{2} + x & , -\pi < x < 0 \\ \frac{\pi}{2} - x & , 0 < x < \pi \end{cases}$ , [8]  
 Hence, deduce that  $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$  and  $\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$
- Q3. a) Find  $L^{-1} \left[ \frac{s^2}{(s^2+1)(s^2+4)} \right]$  using convolution theorem. [6]  
 b) Show that the set of functions  $\phi_n(x) = \sin \left( \frac{n\pi x}{l} \right)$ ,  $n = 1, 2, 3, \dots$  is orthogonal in  $[0, l]$ . [6]  
 c) Using Green's theorem evaluate  $\oint_C (e^{x^2} - xy)dx - (y^2 - ax)dy$  where  $C$  is the circle  $x^2 + y^2 = a^2$ . [8]
- Q4. a) Find Laplace transform of  $f(t) = \begin{cases} \frac{t}{a} & , 0 < t \leq a \\ \frac{(2a-t)}{a} & , a < t < 2a \end{cases}$  and  $f(t) = f(t + 2a)$ . [6]  
 b) Prove that a vector field  $\bar{f}$  is irrotational and hence find its scalar potential  $\bar{f} = (y \sin z - \sin x) i + (x \sin z + 2yz) j + (xy \cos z + y^2) k$ . [6]  
 c) Obtain the Fourier expansion of  $f(x) = \left( \frac{\pi-x}{2} \right)^2$  in the interval  $0 \leq x \leq 2\pi$  and  $f(x + 2\pi) = f(x)$ . Also deduce that  $\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$  [8]
- Q5. a) Use Gauss's Divergence Theorem to evaluate  $\iint_S \bar{N} \cdot \bar{F} ds$  where  $\bar{F} = 4xi + 3yj - 2zk$  and  $S$  is the surface bounded by  $x=0, y=0, z=0$  and  $2x+2y+z=4$ . [6]  
 b) Find the Z-transform of  $f(k) = ke^{-ak}$ ,  $k \geq 0$ . [6]  
 c) i) Find  $L^{-1} \left[ \frac{s+2}{s^2(s+3)} \right]$ . [8]  
 ii) Find  $L^{-1} \left[ \log \left( \frac{s+a}{s+b} \right) \right]$ .
- Q6. a) Solve using Laplace transform  $(D^2 + 3D + 2)y = 2(t^2 + t + 1)$ , with  $y(0) = 2$  and  $y'(0) = 0$ . [6]  
 b) Find the bilinear transformation which maps the points  $Z=1, i, -1$  onto the points  $W=i, 0, -i$ . [6]  
 c) Find Fourier sine integral of  $f(x) = \begin{cases} x & , 0 < x < 1 \\ 2-x & , 1 < x < 2 \\ 0 & , x > 2 \end{cases}$  [8]