



- Note :-**
- 1) Question number 1 is **compulsory**.
  - 2) Attempt any **three** questions from the remaining **five** questions.
  - 3) **Figures** to the **right** indicate **full** marks.

- Q.1
- a) Find the Laplace transform of  $\cos t \cos 2t \cos 3t$ . 05
  - b) Construct an analytic function whose real part is  $e^x \cos y$ . 05
  - c) Find the directional derivative of  $\phi = x^4 + y^4 + z^4$  at point  $A(1, -2, 1)$  in the direction of  $AB$  where  $B$  is  $(2, 6, -1)$ . 05
  - d) Expand  $f(x) = lx - x^2$ ,  $0 < x < l$  in a half-range sine-series. 05
- Q.2
- a) Find the angle between the normals to the surface  $xy = z^2$  at the points  $(1, 4, 2)$ ,  $(-3, -3, 3)$ . 06
  - b) Find the Fourier series for 06
 
$$f(x) = \begin{cases} -c & -a < x < 0 \\ c, & 0 < x < a \end{cases}$$
  - c) Find the inverse Laplace transform of 08
    - (i)  $\frac{4s + 12}{s^2 + 8s + 12}$
    - (ii)  $\log\left(\frac{s^2 + a^2}{\sqrt{s + b}}\right)$
- Q.3
- a) State true or false with proper justification "There does not exist an analytic function whose real part is  $x^3 - 3x^2y - y^3$ ". 06
  - b) Prove that  $J_{5/2}(x) = \sqrt{\frac{2}{\pi x}} \left( \frac{3-x^2}{x^2} \sin x - \frac{3}{x} \cos x \right)$ . 06
  - c) Expand  $f(x) = 4 - x^2$  in the interval  $(0, 2)$ . 08
- Q.4
- a) Use Gauss's Divergence theorem to evaluate  $\iint_S \vec{N} \cdot \vec{F} dS$  where  $\vec{F} = 4x \vec{i} + 3y \vec{j} - 2z \vec{k}$  and  $S$  is the surface bounded by  $x = 0, y = 0, z = 0$  and  $2x + 2y + z = 4$ . 06

TURN OVER

b) Prove that 06  

$$\int x^3 \cdot J_0(x) dx = x^3 \cdot J_1(x) - 2x^2 \cdot J_2(x).$$

c) Solve using Laplace transform  $\frac{dy}{dt} + 3y = 2 + e^{-t}$  with 08  
 $y(0) = 1.$

Q. 5 a) Find Laplace transform of  $(1 + 2t - 3t^2 + 4t^3)H(t - 2)$  where 06  

$$H(t - 2) = \begin{cases} 0, & t < 2 \\ 1, & t \geq 2 \end{cases}$$

b) Prove that  $2J_0''(x) = J_2(x) - J_0(x).$  06

c) Obtain complex form of Fourier Series for  $f(x) = e^{ax}$  in  $(-\pi, \pi)$  08  
 where  $a$  is not an integer. Hence deduce that when  $\alpha$  is a constant other than an integer

$$\sin \alpha x = \frac{\sin \pi \alpha}{i\pi} \sum \frac{(-1)^n n}{(\alpha^2 - n^2)} e^{inx}$$

Q. 6 a) Using Green's theorem evaluate 06

$$\oint_C (e^{x^2} - xy) dx - (y^2 - ax) dy$$

where  $C$  is the circle  $x^2 + y^2 = a^2.$

b) Express the function 06  

$$f(x) = \begin{cases} 1 & \text{for } |x| < 1 \\ 0 & \text{for } |x| > 1 \end{cases}$$
  
 as a Fourier Integral.

c) Under the transformation  $w = (1 + i)z + (2 - i)$ , find the region 08  
 in the  $w$  -plane into which the rectangular region bounded by  
 $x = 0, y = 0, x = 1, y = 2$  in the  $z$  -plane is mapped.

xxx xxx