

Q.P. Code : 39159

(3hours)

[Total marks: 80]



- N.B.** 1) Question No. 1 is compulsory.  
 2) Answer **any Three** from remaining  
 3) Figures to the right indicate full marks

1. a) Find Laplace transform of  $f(t) = e^{-t} \sin t \cdot \cos 2t$ . 5
- b) Show that the set of functions  $\cos nx, n = 1, 2, 3 \dots$  is orthogonal on  $(0, 2\pi)$ . 5
- c) The equations of lines of regression are  $x + 2y = 5$  and  $2x + 3y = -8$ .  
 Find i) means of  $x$  and  $y$ , ii) coefficient of correlation between  $x$  and  $y$ . 5
- d) Evaluate  $\int_C (z^2 - 2\bar{z} + 1) dz$  where  $C$  is the circle  $|z| = 1$ . 5
2. a) Using convolution theorem, find the inverse Laplace transform of 6  

$$F(s) = \frac{1}{(s^2 + 9)(s^2 + 4)}$$
- b) Obtain Fourier series of  $f(x) = |x|$  in  $(-\pi, \pi)$  6
- c) Find the bilinear transformation which maps the points  $z = 1, i, -1$  onto the points  $w = i, 0, -i$ . Hence, find the image of  $|z| < 1$  onto the  $w$ -plane. 8
3. a) If  $v = e^x \sin y$ , prove that  $v$  is a harmonic function. Also find the corresponding harmonic conjugate function and analytic function. 6
- b) Using Bender-Schmidt method, solve  $\frac{\partial^2 u}{\partial x^2} - \frac{\partial u}{\partial t} = 0$ , subject to the conditions,  
 $u(0, t) = 0, u(5, t) = 0, u(x, 0) = x^2(25 - x^2)$  taking  $h = 1$ , for 3 minutes. 6
- c) Using Residue theorem, evaluate
- i)  $\int_0^{2\pi} \frac{d\theta}{2 + \cos\theta}$       ii)  $\int_0^\infty \frac{dx}{x^2 + 1}$  8

[TURN OVER]

4. a) Solve by Crank –Nicholson simplified formula  $\frac{\partial^2 u}{\partial x^2} - \frac{\partial u}{\partial t} = 0$ ,

$u(0, t) = 0$ ,  $u(1, t) = 2t$ ,  $u(x, 0) = 0$  taking  $h = 0.25$  for two-time steps. 6

b) Obtain the Taylor's and Laurent series which represent the function

$$f(z) = \frac{z}{(z-1)(z-2)} \text{ in the regions, i) } |z| < 1 \quad \text{ii) } 1 < |z| < 2 \quad 6$$

c) Solve  $(D^2 - 3D + 2)y = 4e^{2t}$  with  $y(0) = -3$ ,  $y'(0) = 5$  where  $D \equiv \frac{d}{dt}$  8

5. a) Find an analytic function  $f(z) = u + iv$ , if 6

$$u = e^{-x}\{(x^2 - y^2) \cos y + 2xy \sin y\}$$

b) Find the Laplace transform of  $\frac{\sin at}{t}$ . Does the L.T of  $\frac{\cos at}{t}$  exist? 6

c) Obtain half range Fourier cosine series of  $f(x) = x$ ,  $0 < x < 2$ . Using Parseval's identity, deduce that – 8

$$\frac{\pi^4}{96} = \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \dots$$

6. a) Obtain Complex form of Fourier series of  $f(x) = e^x$ ,  $-1 < x < 1$  6

b) Fit a straight line to the following data, 6

$x$	100	120	140	160	180	200
$y$	0.45	0.55	0.60	0.70	0.80	0.85

c) A string is stretched and fastened to two points distance  $l$  apart. Motion is started by displacing the string in form  $y = a \sin(\pi x / l)$  from which it is released at a time  $t = 0$ . If the vibrations of a string is given by  $\frac{\partial^2 y}{\partial t^2} = c^2 \frac{\partial^2 y}{\partial x^2}$ , show that the displacement of a point at a distance  $x$  from one end at time  $t$  is given by  $y(x, t) = a \sin(\pi x / l) \cos(\pi ct / l)$ . 8