

(3 Hours)

[Total Marks: 80]

- N.B.: (1) Question No.1 is compulsory.
(2) Answer any **three** from remaining **five** questions.
(3) Figures to the **right** indicate **full** marks.
(4) Assume the **data** if it is **necessary**.
(5) **Vector notation** must be used wherever necessary.

Q.1) Attempt any **four** of the following:- (05-Marks each)

[20]

- (a) Find the charge enclosed in a cube of having side of 2 m with the edges of the cube parallel to axes x, y, and z while origin is at the centre of the cube. The charge density within the cube is $50x^2 \cos\left(\frac{\pi}{2}y\right) \mu\text{C}/\text{m}^3$.
- (b) Explain the concept of potential gradient and the relation between electric field and potential.
- (c) If the magnetic field $\vec{H} = (3x \cos \beta + 6z \sin \alpha) \hat{a}_y$. Find the current density \vec{J} if field are invariant with time.
- (d) Discuss the phenomenon of polarization in dielectric medium. Also discuss how it gives rise to bound charge densities.
- (e) For a lossy dielectric material having $\mu_r = 1$, $\epsilon_r = 48$ and $\sigma = 20 \text{ s/m}$. Calculate the propagation constant at a frequency of 16 GHz.

Q.2)

[20]

- (a) Given $\vec{D} = 2rz \cos^2 \phi \hat{a}_r - rz \sin \phi \cos \phi \hat{a}_\phi + r^2 \cos^2 \phi \hat{a}_z$. Calculate electric flux through the following surfaces.
(i) $r = 3$, $0 \leq z \leq 5$. (ii) $z = 0$, $0 \leq r \leq 3$.
- (b) Obtain E inside, outside solid sphere. A uniform volume charge density $\rho_v \text{ C}/\text{m}^3$, Distributed in a solid sphere of radius 'a' find expression of E everywhere.

[10]

[10]

Q.3)

[20]

- (a) Planes $z = 0$ and $z = 4$ carry a current $\vec{K} = -10 \hat{a}_x \text{ A/m}$ and $\vec{K} = 10 \hat{a}_x \text{ A/m}$ respectively. Find \vec{H} at points (i) $P(1, 1, 1)$ and (ii) $Q(0, -3, 10)$
- (b) Obtain an expression for magnetic vector potential in the region surrounding an infinitely long straight filamentary current 'I'.

[10]

[10]

Q.4)

[20]

- (a) Derive the Poisson's and Laplace equation. And the one dimensional Laplace's equation is as $\frac{\partial^2 V}{\partial x^2} = 0$, The boundary conditions are $V = 9$ at $X = 1$ and $V = 0$ at $X = 10$. Find the potential and show the variation of V with respect to X.

[10]

[TURN OVER

- (b) A potential field is given as $V = 100 e^{-5x} \sin 3y \cos 4z$ volts. Let point $P(0.1, \frac{\pi}{12}, \frac{\pi}{24})$ be located at a conductor free space boundary. At point P, find the magnitudes of (i) V , (ii) \vec{E} , (iii) E_t , (iv) E_N , (v) \vec{D} , (vi) D_N , and (vii) ρ_s . [10]

Q.5)

- (a) Derive the set of Maxwell's equations for static fields and harmonically time varying fields. [20]

- (b) Verify whether the following fields [10]

$$\vec{E} = (2 \cos x \sin t) \hat{a}_y \text{ and } \vec{H} = \left(\frac{2}{\mu_0} \cos x \cos t\right) \hat{a}_z. \text{ Satisfy Maxwell's equation in free space. [10]}$$

Q.6)

- (a) Formulate the wave equation from Maxwell's equations. Solve it for perfectly conducting media. [20]

- (b) The magnetic field intensity of a uniform plane wave in air is 20 A/m along the \hat{a}_y direction. The wave is propagating in the \hat{a}_z direction at a frequency of $2 \times 10^9 \text{ rad/sec}$. Find the [10]

- (i) Wavelength, (ii) Frequency, (iii) Period, and (iv) Amplitude of \vec{E} .