

**MATHEMATICS SOLUTION**  
**(DEC 2019 SEM 4 MECHANICAL)**

**Q1) (a) Find eigen values of  $A^2 - 2A + I$  and  $\text{adj } A$  Where  $A = \begin{bmatrix} 4 & 1 & -1 \\ 6 & 3 & -5 \\ 6 & 2 & -2 \end{bmatrix}$ . (5M)**

**Solution:**

$$A = \begin{bmatrix} 4 & 1 & -1 \\ 6 & 3 & -5 \\ 6 & 2 & -2 \end{bmatrix}$$

$$\begin{aligned} A^2 - 2A + I &= \begin{bmatrix} 4 & 1 & -1 \\ 6 & 3 & -5 \\ 6 & 2 & -2 \end{bmatrix} \begin{bmatrix} 4 & 1 & -1 \\ 6 & 3 & -5 \\ 6 & 2 & -2 \end{bmatrix} - 2 \begin{bmatrix} 4 & 1 & -1 \\ 6 & 3 & -5 \\ 6 & 2 & -2 \end{bmatrix} + \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 16 & 5 & -7 \\ 12 & 5 & -11 \\ 24 & 8 & -12 \end{bmatrix} - \begin{bmatrix} 8 & 2 & -2 \\ 12 & 6 & -10 \\ 12 & 4 & -4 \end{bmatrix} + \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 9 & 3 & -5 \\ 0 & 0 & -1 \\ 12 & 4 & -7 \end{bmatrix} \end{aligned}$$

The Characteristic equation is

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} 9 - \lambda & 3 & -5 \\ 0 & 0 - \lambda & -1 \\ 12 & 4 & -7 - \lambda \end{vmatrix} = 0$$

$$(9 - \lambda)[(-\lambda)(-7 - \lambda) + 4] - 3[12] - 5[12\lambda] = 0$$

$$-\lambda(\lambda^2 - 2\lambda + 1) = 0$$

$$-\lambda(\lambda - 1)^2 = 0$$

$$\lambda = 0, 1, 1$$

Hence the eigen values of  $A^2 - 2A + I$  are 0, 1 and 1.

**(b) A random variable X has the following probability function.**

<b>X</b>	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
<b>P(X=x)</b>	<b>1/16</b>	<b>4k</b>	<b>6k</b>	<b>4k</b>	<b>k</b>

**Find (i) k (ii) P (X < 4) (iii) P (X > 3) (iv) P (0 < X ≤ 2). (5M)**

**Solution:**

(i) Since  $\sum p(x_i) = 1$

$$1/16 + 4k + 6k + 4k + k = 1$$

$$1/16 + 15k = 1$$

$$15k = 1 - 1/16$$

$$k = 1/16$$

Thus, we have the following probability distribution

X	0	1	2	3	4
P(X=x)	1/16	4/16	6/16	4/16	1/16

$$(ii) P(X < 4) = P(X = 0, 1, 2, 3)$$

$$= P(X = 0) + P(X = 1) + P(X = 2) + P(X = 3)$$

$$= 1/16 + 4/16 + 6/16 + 4/16$$

$$= 15/16.$$

$$(iii) P(X > 3) = P(X = 4)$$

$$= 1/16.$$

$$(iv) P(0 < X \leq 2) = P(X = 1) + P(X = 2)$$

$$= 4/16 + 6/16$$

$$= 10/16.$$

**(c) Can it be concluded that the average life span of an Indian is more than 71 years, if a random sample of 900 Indians has an average life span 72.8 years with standard deviation of 7.2 years? (5M)**

**Solution:**

Null Hypothesis  $H_0$ :  $\mu = 70$  years

Alternate Hypothesis  $H_a$ :  $\mu \neq 70$  years

$$\text{Test statistic: } Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$$

Since we are given standard deviation of the sample, we put

$$\bar{X} = 71.8, \mu = 70, \sigma = 7.2, n = 100$$

$$Z = \frac{71.8 - 70}{7.2/\sqrt{100}} = 2.5$$

Level of significance:  $\alpha = 0.05$

Critical value: the value of  $z_\alpha$  at 5% level of significance is 1.96

Decision: Since the computed value  $|Z| = 2.5$  is greater than the critical value  $z_\alpha = 1.96$ , the null hypothesis is rejected.

(d) Consider the following problem

$$\text{Maximize } Z = 2x_1 - 2x_2 + 4x_3 - 5x_4$$

$$\text{Subjected to } x_1 + 4x_2 - 2x_3 + 8x_4 = 2$$

$$-x_1 + 2x_2 + 3x_3 + 4x_4 = 1$$

$$x_1, x_2, x_3, x_4 \geq 0$$

Find a basic feasible solution which is non-degenerate and optimal solution.

Solution:

No. of basic solutions	Non-basic variables = 0	Basic variables	Equations and values of the basic variables	Is the solution feasible?	Is the solution degenerate?	Value Of z	Is the solution optimal?
1	$x_3=0$ $x_4=0$	$x_1, x_2$	$x_1 + 4x_2 = 2$ $-x_1 + 2x_2 = 1$ $x_1 = 0, x_2 = 1/2$	Yes	Yes	-1.5	No
2	$x_2=0$ $x_4=0$	$x_1, x_3$ outgoing $x_2$ incoming $x_1$	$x_1 - 2x_3 = 2$ $-x_1 + 3x_3 = 1$ $x_1 = 8, x_3 = 3$	Yes	No	28	Yes
3	$x_1=0$ $x_4=0$	$x_2, x_3$ outgoing $x_1$ incoming $x_2$	$4x_2 - 2x_3 = 2$ $2x_2 + 3x_3 = 1$ $x_3 = 0, x_2 = 1/2$	Yes	Yes	-1	No
4	$x_2=0$ $x_3=0$	$x_1, x_4$ outgoing $x_2$ incoming $x_1$	$x_1 + 8x_4 = 2$ $-x_1 + 4x_4 = 1$ $x_1 = 0, x_4 = 1/4$	Yes	Yes	-1.25	No
5	$x_1=0$ $x_3=0$	$x_2, x_4$ outgoing $x_1$ incoming $x_2$	$4x_2 + 8x_4 = 2$ $2x_2 + 4x_4 = 1$ Unbounded	---	---	---	---
6	$x_1=0$ $x_2=0$	$x_3, x_4$ outgoing $x_2$ incoming $x_3$	$-2x_3 + 8x_4 = 2$ $3x_3 + x_4 = 12$ $x_3 = 0, x_4 = 1/4$	Yes	Yes	-12.5	No

Q2(a) Check whether the given matrix A is diagonalizable, diagonalize if it is, where  $A = \begin{bmatrix} 4 & 1 & -1 \\ 6 & 3 & -5 \\ 6 & 2 & -2 \end{bmatrix}$

Solution:

(6M)

The Characteristic equation is

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} 4 - \lambda & 1 & -1 \\ 6 & 3 - \lambda & -5 \\ 6 & 2 & -2 - \lambda \end{vmatrix} = 0$$

$$-\lambda^3 + 5\lambda^2 - 8\lambda + 4 = 0$$

$$-(\lambda - 1)(\lambda^2 - 4\lambda + 4) = 0$$

$$-(\lambda - 1)(\lambda - 2)(\lambda - 2) = 0$$

[MUQuestionPapers.com](http://MUQuestionPapers.com)

$$\lambda = 1, 2, 2.$$

When  $\lambda = 1$ ,

$$\begin{bmatrix} 3 & 1 & -1 \\ 6 & 2 & -5 \\ 6 & 2 & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

By  $R_1/3$

$$\begin{bmatrix} 1 & 1/3 & -1/3 \\ 6 & 2 & -5 \\ 6 & 2 & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

By  $R_2 - 6R_1$

$$\begin{bmatrix} 1 & 1/3 & -1/3 \\ 0 & 0 & 3 \\ 6 & 2 & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

By  $R_3 - 6R_1$

$$\begin{bmatrix} 1 & 1/3 & -1/3 \\ 0 & 0 & -3 \\ 0 & 0 & -1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

By  $R_2/-3$

$$\begin{bmatrix} 1 & 1/3 & -1/3 \\ 0 & 0 & 1 \\ 0 & 0 & -1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

By  $R_3 - (-1)R_2$

$$\begin{bmatrix} 1 & 1/3 & -1/3 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

By  $R_1 - (-1/3)R_2$

$$\begin{bmatrix} 1 & 1/3 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$x_3 = 0$$

$$x_1 + \left(\frac{1}{3}\right)x_2 = 0$$

Therefore,

$$x_1 = \left(-\frac{1}{3}\right)x_2$$

$$x_2 = x_2$$

$$x_3 = 0$$

The rank of coefficient matrix is 2. The number of unknowns is 3. Hence, there are  $3-2 = 1$ . Putting  $x_2 = t$  then  $x_1 = \left(-\frac{t}{3}\right)$ .

$$X_1 = \begin{bmatrix} -t/3 \\ t \\ 0 \end{bmatrix} = t \begin{bmatrix} -1/3 \\ 1 \\ 0 \end{bmatrix}$$

Corresponding to  $\lambda = 1$ , we get the eigenvector  $X_1' = [-1/3 \quad 1 \quad 0]$

There are three variables and the rank is 2, hence there is only one independent solution.

For  $\lambda = 1$ , algebraic multiplicity is 1 and geometric multiplicity is 1.

When  $\lambda = 2$ ,

$$\begin{bmatrix} 2 & 1 & -1 \\ 6 & 1 & -5 \\ 6 & 2 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

By  $R_1/2$

$$\begin{bmatrix} 1 & 1/2 & -1/2 \\ 6 & 1 & -5 \\ 6 & 2 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

By  $R_2 - 6R_1$

$$\begin{bmatrix} 1 & 1/2 & -1/2 \\ 0 & -2 & -2 \\ 6 & 2 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

By  $R_3 - 6R_1$

$$\begin{bmatrix} 1 & 1/2 & -1/2 \\ 0 & -2 & -2 \\ 0 & -1 & -1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

By  $R_2/-2$

$$\begin{bmatrix} 1 & 1/2 & -1/2 \\ 0 & 1 & 1 \\ 0 & -1 & -1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

By  $R_3 - (-1)R_2$

$$\begin{bmatrix} 1 & 1/2 & -1/2 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

By  $R_1 - (-1/2)R_2$

$$\begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$x_1 - x_3 = 0$$

$$x_2 + x_3 = 0$$

Therefore,

$$x_1 = x_3$$

$$x_2 = -x_3$$

$$x_3 = x_3$$

The rank of coefficient matrix is 2. The number of unknowns is 3. Hence, there are  $3-2 = 1$ . Putting  $x_3 = t$

$$X_1 = \begin{bmatrix} t \\ -t \\ t \end{bmatrix} = t \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

There are three variables and the rank is 2, hence there is only one independent solution.

For  $\lambda = 2$ , algebraic multiplicity is 2 and geometric multiplicity is 1.

Hence the algebraic multiplicity and geometric multiplicity should be same.

Thus, the matrix is not diagonalized.

**(b) Verify Green's theorem for  $\bar{F} = x^2i - xyj$  where C is the triangle having vertices A (0,3), B (3,0), C (6,3). (6M)**

**Solution:**

By Green's Theorem

$$\int_C P \cdot dx + Q \cdot dy = \iint_R \left( \frac{\delta Q}{\delta x} - \frac{\delta P}{\delta y} \right) \cdot dx \cdot dy$$

Here,  $P = x^2$  ;  $Q = -xy$

$$\frac{\delta Q}{\delta x} = -y ; \frac{\delta P}{\delta y} = 0$$

(a) Along AB, since the equation of AB is

$$\frac{y-3}{3-0} = \frac{x-0}{0-3}$$

$$y = 3 - x$$

Putting  $P = x^2$  ;  $Q = -xy = -x(3 - x)$ ;  $dy = -dx$

$$\begin{aligned} \int_C P \cdot dx + Q \cdot dy &= \int_0^3 [x^2 + x(3 - x)] dx = \int_0^3 3x \cdot dx \\ &= (3x^2/2 | x = 0 \text{ to } 3) \\ &= \frac{27}{2} \end{aligned}$$

Along BC, since the equation of BC,  $\frac{y-0}{0-3} = \frac{x-3}{3-6}$  i.e., is  $y = x - 3$

$$\begin{aligned} \int_C P \cdot dx + Q \cdot dy &= \int_3^6 (x^2 - x^2 + 3x) dx = \int_3^6 3x \cdot dx \\ &= (3x^2/2 | x = 3 \text{ to } 6) \\ &= \frac{81}{4} \end{aligned}$$

Along CA, since the equation of CA is  $y = 3$  ;  $dy = 0$

$$\begin{aligned} \int_C P \cdot dx + Q \cdot dy &= \int_6^0 (x^2 - x^2 + 3x) dx = \int_6^0 x^2 \cdot dx \\ &= (x^3/3 | x = 6 \text{ to } 0) \\ &= -54 \end{aligned}$$

(b)

$$\begin{aligned} \iint_R \left( \frac{\delta Q}{\delta x} - \frac{\delta P}{\delta y} \right) \cdot dx \cdot dy &= \int_0^3 \int_{3-y}^{3+y} (-y) \cdot dx \cdot dy \\ &= \int_0^3 (-y \cdot x | y = 3 - y \text{ to } 3 + y) \cdot dx \end{aligned}$$

$$\begin{aligned}
&= \int_0^3 y[3 + y - 3 + y] \cdot dx \\
&= \int_0^1 y^3 \cdot dx \\
&= \left(\frac{y^4}{4} \Big|_{x=0}^3\right) \\
&= \frac{81}{4}
\end{aligned}$$

From (a) and (b), the theorem is verified.

**(c) Sample of two types of two types of electric bulbs were tested for length of life and the following data were obtained**

	Type I	Type II
Sample size	10	9
Mean of the sample (in hours)	1136	1034
Standard deviation	36	39

**Test at 5% level of significance whether the difference in the sample means is significant. (8M)**

**Solution:**

Null Hypothesis  $H_0$ : There is no relation between the electric bulbs and length of life.

Alternate Hypothesis  $H_a$ : There is a relation between these two.

To test the significant difference between two mean  $\bar{x}_1$  and  $\bar{x}_2$  of sample sizes  $n_1$  and  $n_2$  use the statistic

$$t = \frac{(\bar{x}_1 - \bar{x}_2)}{s \sqrt{\left(\frac{1}{n_1}\right) + \left(\frac{1}{n_2}\right)}}$$

$$s^2 = \frac{(n_1 s_1^2 + n_2 s_2^2)}{n_1 + n_2 - 2} = \frac{(8 \cdot 36 \cdot 36 + 7 \cdot 40 \cdot 40)}{8 + 7 - 2}$$

$$s = 40.7$$

$$t = \frac{(1234 - 1036)}{40.7 \cdot \sqrt{\frac{1}{8} + \frac{1}{7}}} = 9.39$$

Calculated t value = 9.39

Tabulated Value = 1.77 (as 5% level of significance with 13 degrees of freedom)

$9.39 > 1.77$ , reject  $H_0$  (Null Hypothesis)

Hence, there is no relation between the electric bulbs and length of life.

**Q3 (a) Use the dual simplex method to solve the following LPP**

**Minimise  $Z = 6x_1 + x_2$**

**Subject to  $2x_1 + x_2 \geq 3$**

$x_1 - x_2 \geq 0$

$x_1, x_2 \geq 0.$

(6M)

**Solution:**

Minimise  $z = 6x_1 + x_2$

Subject to  $-2x_1 - x_2 \leq -3$

$-x_1 + x_2 \leq 0$

Introducing the slack variables  $s_1, s_2$ , we have

Minimise  $z = 6x_1 + x_2 - 0s_1 - 0s_2$

i.e.  $z - 6x_1 - x_2 + 0s_1 + 0s_2$

Subject to  $-2x_1 - x_2 + s_1 + 0s_2 = -3$

$-x_1 + x_2 + 0s_1 + s_2 = 0$

Iteration Number	Basic Variable	Coefficient Of				R.H.S. Solution
		$x_1$	$x_2$	$s_1$	$s_2$	
0	z	-6	-1	0	0	0
$s_1$ leaves	$s_1$	-2	-1*	1	0	-3
$x_2$ enters	$s_2$	-1	1	0	1	0
Ratio		3	1			
1	z	-4	0	-1	0	3
$s_2$ leaves	$x_2$	2	1	-1	0	3
$x_1$ leaves	$s_2$	-3*	0	1	1	-3
2	z	0	0	-7/3	4/3	7
	$x_2$	0	1	-1/3	2/3	1
	$x_1$	1	0	-1/3	-1/3	1

$x_1 = 1, x_2 = 1$  and  $z_{\min} = 7$

**(b) Use Gauss Divergence Theorem to evaluate  $\iint_S \bar{N} \cdot \bar{F} \cdot d\bar{s}$  where  $\bar{F} = 2xi + 2yj + 2z^2k$  and S is the closed surface bounded by the cone  $x^2 + y^2 = z^2$  and the plane  $z = 1$ .** (6M)

**Solution:**

By divergence formula,

$\iint_S \bar{F} \cdot d\bar{S} = \iiint_V \nabla \cdot \bar{F} \cdot div$

Now,  $\bar{F} = 2xi + 2yj + 2z^2k$



$$\begin{aligned}\nabla \cdot \bar{F} &= \left( \frac{\delta(2x)}{\delta x} + \frac{\delta(2y)}{\delta y} + \frac{\delta(2z^2)}{\delta z} \right) \\ &= 2 + 2 + 2z \\ &= 4 + 2z \\ &= 2(2 + z)\end{aligned}$$

$$\iiint_V \nabla \cdot \bar{F} \cdot \text{div} = \iiint_V 2(2 + z) \cdot \text{div} = \iiint_V 2(2 + z) \cdot dx \cdot dy \cdot dz$$

We shall obtain the volume integral by using cylinder co-ordinates  $x = r \cdot \cos\theta$ ,  $y = r \cdot \sin\theta$ ,  $z = z$  and  $dx \cdot dy \cdot dz = r \cdot dr \cdot d\theta \cdot dz$

$$r^2 = x^2 + y^2 \text{ and by data } x^2 + y^2 = z^2$$

$z = r$ , Hence,  $z$  varies from 0 to 1

$$\begin{aligned}\iiint_V 2(2 + z) \cdot dx \cdot dy \cdot dz &= \int_{\theta=0}^{2\pi} \int_{r=0}^1 \int_{z=r}^1 2(2 + z) \cdot r \cdot dr \cdot d\theta \cdot dz \\ &= 2 \int_0^{2\pi} \int_0^1 \left( 2rz + r \frac{z^2}{2} \Big|_{z=r \text{ to } 1} \right) \cdot dr \cdot d\theta \\ &= 2 \int_0^{2\pi} \int_0^1 \left( \frac{5r}{2} - 2r^2 - \frac{r^3}{2} \right) \cdot dr \cdot d\theta \\ &= 2 \int_0^{2\pi} \left( \frac{5r^2}{4} - \frac{2r^3}{3} + \frac{r^4}{8} \Big|_{r=0 \text{ to } 1} \right) \cdot d\theta \\ &= 2 \int_0^{2\pi} \left( \frac{5}{4} - \frac{2}{3} + \frac{1}{8} \right) \cdot d\theta \\ &= 2 \int_0^{2\pi} \frac{17}{24} \cdot d\theta \\ &= \left( \frac{17}{12} \theta \Big|_{\theta=0 \text{ to } 2\pi} \right) \\ &= \frac{17}{12} * 2\pi = \frac{17\pi}{6}\end{aligned}$$

(c) Find the rank, index, signature and class of the following Quadratic form by reducing it to its canonical form

$$2x^2 - 2y^2 + 2z^2 - 2xy - 8yz + 6zx$$

(8M)

**Solution:**

The quadratic form can be written as

The matrix form is

$$A = \begin{bmatrix} 2 & -1 & 3 \\ -1 & 2 & -4 \\ 3 & -4 & 2 \end{bmatrix}$$

We write  $A = IAI$

$$\begin{bmatrix} 2 & -1 & 3 \\ -1 & 2 & -4 \\ 3 & -4 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} A \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{By } R_2 + \frac{1}{2}R_1, R_3 - \frac{3}{2}R_1, C_2 + \frac{1}{2}C_1, C_3 - \frac{3}{2}C_1$$

$$\begin{bmatrix} 2 & 0 & 0 \\ 0 & -5/2 & -5/2 \\ 0 & -5/2 & -5/2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 1/2 & 1 & 0 \\ -3/2 & 0 & 1 \end{bmatrix} A \begin{bmatrix} 1 & 1/2 & -3/2 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{By } R_3 - R_2, C_3 - C_2$$

$$\begin{bmatrix} 2 & 0 & 0 \\ 0 & -5/2 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 1/2 & 1 & 0 \\ -2 & -1 & 1 \end{bmatrix} A \begin{bmatrix} 1 & 1/2 & -2 \\ 0 & -1 & -1 \\ 0 & 0 & 1 \end{bmatrix}$$

By  $\frac{1}{\sqrt{2}}R_1, \sqrt{\frac{2}{5}}R_2, \frac{1}{\sqrt{2}}C_1, \sqrt{\frac{2}{5}}C_2$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 1/\sqrt{2} & 0 & 0 \\ 1/\sqrt{10} & \sqrt{2/5} & 0 \\ -2 & -1 & 1 \end{bmatrix} A \begin{bmatrix} 1/\sqrt{2} & 1/\sqrt{10} & -2 \\ 0 & \sqrt{2/5} & -1 \\ 0 & 0 & 1 \end{bmatrix}$$

The linear transform,  $X = PY$

$$x = \frac{1}{\sqrt{2}}u + \frac{1}{\sqrt{10}}v - 2w$$

$$y = -\sqrt{\frac{2}{5}}u - w$$

$$z = w$$

Transforms the given quadratic form  $u^2 + v^2$

The rank = 2, Index = 1, Signature = 1

The form is positive semi-definite.

**Q4 (a) Four dice were thrown 250 times and the number of appearance of 6 each time was noted**

No. of successes(x):	0	1	2	3	4
Frequency (f):	133	69	34	11	3

Fit a poisson distribution and find the expected frequencies for  $x = 1, 2, 3, 4$ .

(6M)

**Solution:**

$$\text{Now, mean} = m = \frac{\sum f_i x_i}{\sum f_i}$$

$$\text{Mean} = \frac{133 \cdot 0 + 69 \cdot 1 + 34 \cdot 2 + 11 \cdot 3 + 3 \cdot 4}{250} = 0.728$$

Poisson distribution of X is

$$P(X = x) = \frac{e^{-m} X m^x}{x!} = \frac{e^{-0.728} X 0.728^x}{x!}$$

Expected frequency =  $N \times p(x)$

$$= 250 X \frac{e^{-0.728} X 0.728^x}{x!}$$

Putting  $x = 0, 1, 2, 3, 4$  we get the expected frequencies as 133, 69, 34, 11, 3.

$$f(x+1) = \frac{m}{x+1} \cdot f(x) = \frac{0.728}{x+1} \cdot f(x)$$

$$\text{Putting } x = 0, f(1) = \frac{0.728}{0+1} * 133 = 96.44$$

$$\text{Putting } x = 1, f(2) = \frac{0.728}{1+1} * 69 = 25.26$$

$$\text{Putting } x = 2, f(3) = \frac{0.728}{2+1} * 34 = 8.11$$

$$\text{Putting } x = 3, f(4) = \frac{0.728}{3+1} * 11 = 2.02$$

(b) Verify Cayley Hamilton theorem for matrix A and hence find the matrix represented by  $A^5 - 4A^4 - 7A^3 + 11A^2 - A - 11I$

$$\text{Where } A = \begin{bmatrix} 3 & -2 & 3 \\ 10 & -3 & 5 \\ 5 & -4 & 7 \end{bmatrix}.$$

(6M)

**Solution:**

The characteristic equation is

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} 3 - \lambda & -2 & 3 \\ 10 & -3 - \lambda & 5 \\ 5 & -4 & 7 - \lambda \end{vmatrix} = 0$$

$$-\lambda^3 + 7\lambda^2 - 16\lambda + 12 = 0$$

By Cayley Hamilton theorem, this equation is satisfied by A

$$-A^3 + 7A^2 - 16A + 12I = 0$$

$$A^2 = \begin{bmatrix} 3 & -2 & 3 \\ 10 & -3 & 5 \\ 5 & -4 & 7 \end{bmatrix} \begin{bmatrix} 3 & -2 & 3 \\ 10 & -3 & 5 \\ 5 & -4 & 7 \end{bmatrix} = \begin{bmatrix} 4 & -12 & 20 \\ 25 & -31 & 50 \\ 10 & -26 & 44 \end{bmatrix}$$

$$A^3 = \begin{bmatrix} 4 & -12 & 20 \\ 25 & -31 & 50 \\ 10 & -26 & 44 \end{bmatrix} \begin{bmatrix} 3 & -2 & 3 \\ 10 & -3 & 5 \\ 5 & -4 & 7 \end{bmatrix} = \begin{bmatrix} -8 & -52 & 92 \\ 15 & -157 & 270 \\ -10 & -118 & 208 \end{bmatrix}$$

$$\begin{aligned} -A^3 + 7A^2 - 16A + 12I &= - \begin{bmatrix} -8 & -52 & 92 \\ 15 & -157 & 270 \\ -10 & -118 & 208 \end{bmatrix} + 7 \begin{bmatrix} 4 & -12 & 20 \\ 25 & -31 & 50 \\ 10 & -26 & 44 \end{bmatrix} - 16 \begin{bmatrix} 3 & -2 & 3 \\ 10 & -3 & 5 \\ 5 & -4 & 7 \end{bmatrix} + 12 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \end{aligned}$$

(c) An investigation into the equality of standard deviation of two normal populations gave the following results.

Sample	Size	Sample mean	Sum of squares of deviations from the mean
1	13	18	105
2	21	24	145

Examine the equality of sample variances at 5% level of significance.

(Given:  $F_{0.025} = 2.68$  for d.o.f. 12 and 20 and  $F_{0.025} = 3.07$  for d.o.f. 20 and 12)

(8M)

**Solution:**

Null Hypothesis  $H_0: \sigma_1^2 = \sigma_2^2$

Alternative Hypothesis  $H_a: \sigma_1^2 \neq \sigma_2^2$

Calculations of Test Statistic:  $F = \frac{n_1 s_1^2 / (n_1 - 1)}{n_2 s_2^2 / (n_2 - 1)}$

But  $n_1 s_1^2 = \sum(x_i - \bar{x})^2$  and  $n_2 s_2^2 = \sum(y_i - \bar{y})^2$

$$F = \frac{105/12}{145/20} = \frac{8.75}{7.25} = 1.207$$

Level of significance  $\alpha=0.05$

Degree of freedom  $v_1 = n_1 - 1 = 12$  for the numerator

$v_2 = n_2 - 1 = 20$  for the denominator

Critical Value: The table value

$$F_{(12,20)}(0.025)=2.68$$

$$F_{(20,12)}(0.025)=3.07$$

$$\frac{1}{F_{(20,12)}(0.025)} = \frac{1}{3.07} = 0.326$$

Decision: Since the calculated value  $F=1.207$  lies between 0.326 and 3.07, we accept the null hypothesis.

**Q5 (a) Is matrix  $A = \begin{bmatrix} 2 & 0 & 0 \\ -3 & 3 & -1 \\ 3 & -1 & 3 \end{bmatrix}$  Derogatory matrix? Find its minimal polynomial. (6M)**

**Solution:**

The Characteristic equation is

$$|A - \lambda I| = 0$$

$$\begin{vmatrix} 2 - \lambda & 0 & 0 \\ -3 & 3 - \lambda & -1 \\ 3 & -1 & 3 - \lambda \end{vmatrix} = 0$$

$$(2 - \lambda)[(3 - \lambda)(3 - \lambda) - 1] = 0$$

$$-(2 - \lambda)(\lambda^2 - 6\lambda + 8) = 0$$

$$-(2 - \lambda)(2 - \lambda)(4 - \lambda) = 0$$

$$\lambda = 2, 2, 4.$$

Let us find the minimal polynomial of A. We know that each characteristic root of A is also a root of the minimal polynomial of A. So if  $f(x)$  is the minimal polynomial of A, then  $x - 2$  and  $x - 4$  are the factors of  $f(x)$ . Let us see whether  $(x - 2)(x - 4) = x^2 - 6x + 8$  annihilates A.

Now,  $A^2 - 6A + 8I$

$$= \begin{bmatrix} 2 & 0 & 0 \\ -3 & 3 & -1 \\ 3 & -1 & 3 \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 \\ -3 & 3 & -1 \\ 3 & -1 & 3 \end{bmatrix} - 6 \begin{bmatrix} 2 & 0 & 0 \\ -3 & 3 & -1 \\ 3 & -1 & 3 \end{bmatrix} + 8 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 4 & -18 & 18 \\ 0 & 10 & -6 \\ 0 & -6 & 10 \end{bmatrix} - 6 \begin{bmatrix} 2 & 0 & 0 \\ -3 & 3 & -1 \\ 3 & -1 & 3 \end{bmatrix} + 8 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$f(x) = x^2 - 6x + 8$  annihilates A.

Thus,  $f(x)$  is the monic polynomial of the lowest degree that annihilates A.

Hence,  $f(x)$  is the minimal polynomial of  $A$ . Since the degree of  $f(x)$  is less than the order of  $A$ .  $A$  is derogatory.

(b) A vector field  $\bar{F}$  is given by

$$\bar{F} = (y \sin x - \sin x)\mathbf{i} + (x \sin z + 2yz)\mathbf{j} + (xy \cos z + y^2)\mathbf{k}$$

Prove that  $\bar{F}$  is irrotational. Hence find its scalar potential function  $\phi$  and  $\phi(\pi, 1, 0)$ . (6M)

**Solution:**

We have,

$$\begin{aligned} \text{Curl } \bar{F} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y \sin x - \sin x & x \sin z + 2yz & xy \cos z + y^2 \end{vmatrix} \\ &= \mathbf{i} \left( \frac{\partial}{\partial y} (xy \cos z + y^2) - \frac{\partial}{\partial z} (x \sin z + 2yz) \right) - \mathbf{j} \left( \frac{\partial}{\partial x} (xy \cos z + y^2) - \frac{\partial}{\partial z} (y \sin x - \sin x) \right) \\ &\quad + \mathbf{k} \left( \frac{\partial}{\partial x} (x \sin z + 2yz) - \frac{\partial}{\partial y} (y \sin x - \sin x) \right) \\ &= [x \cos z + 2y - x \cos z - 2y]\mathbf{i} + [y \cos z - y \cos z]\mathbf{j} + [\sin z - \sin x]\mathbf{k} \\ &= 0\mathbf{i} + 0\mathbf{j} + 0\mathbf{k} \end{aligned}$$

Hence,  $\bar{F}$  is irrotational.

If  $\phi$  is the scalar potential then  $\bar{F} = \nabla \phi$

$$(y \sin x - \sin x)\mathbf{i} + (x \sin z + 2yz)\mathbf{j} + (xy \cos z + y^2)\mathbf{k} = \frac{\partial \phi}{\partial x} \mathbf{i} + \frac{\partial \phi}{\partial y} \mathbf{j} + \frac{\partial \phi}{\partial z} \mathbf{k}$$

$$\frac{\partial \phi}{\partial x} = (y \sin x - \sin x) \dots \dots \dots \text{(i)}$$

$$\frac{\partial \phi}{\partial y} = (x \sin z + 2yz) \dots \dots \dots \text{(ii)}$$

$$\frac{\partial \phi}{\partial z} = (xy \cos z + y^2) \dots \dots \dots \text{(iii)}$$

$$\begin{aligned} \text{But, } d\phi &= \frac{\partial \phi}{\partial x} dx + \frac{\partial \phi}{\partial y} dy + \frac{\partial \phi}{\partial z} dz \\ &= (y \sin x - \sin x)dx + (x \sin z + 2yz)dy + (xy \cos z + y^2)dz \\ &= [y \sin z dx + x \sin z dy + xy \cos z dz] + (-\sin x)dx + (2yz dy + y^2 dz) \end{aligned}$$

By integration,  $\phi = x y \sin z + \cos x + y^2 z + c$  where  $c$  is the constant of integration.

Hence putting  $x = \pi$ ,  $y = 1$  and  $z = 0$

$$\begin{aligned} \text{Now, } \phi(\pi, 1, 0) &= \pi * 1 * \sin 0 + \cos \pi + (1)^2 * 0 + c \\ &= \cos \pi + c \end{aligned}$$

(c) The following table gives the result of opinion pole for three vehicles A, B, C. Test whether the age and the choice of the vehicle are independent at 5% level of significance using chi – test.

Age	Vehicle			Total
	A	B	C	
20-35	25	40	35	100
35-50	35	24	41	100
Above 50	40	36	24	100
<b>Total</b>	<b>100</b>	<b>100</b>	<b>100</b>	<b>300</b>

**Solution:**

(8M)

Null Hypothesis  $H_0$ : There is no relation between the age and choice of vehicle.

Alternate Hypothesis  $H_a$ : There is a relation between these two.

On the basis on this hypothesis the number in the first cell =  $\frac{A \times B}{N}$

where, A = total in the first column

B = total in the first row,

N = total number of observations

The number in the first cell of the first row =  $\frac{100 \times 100}{300} = 33.33$

Similarly, The number in the second cell of the first row =  $\frac{100 \times 100}{300} = 33.33$

The number in the first cell of the second row =  $\frac{100 \times 100}{300} = 33.33$

Since the total remain the same, the numbers in the remaining cells are 33.33

Thus, we get the following table

Age	Vehicle			Total
	A	B	C	
20-35	33.33	33.33	33.33	100
35-50	33.33	33.33	33.33	100
Above 50	33.33	33.33	33.33	100
<b>Total</b>	<b>100</b>	<b>100</b>	<b>100</b>	<b>300</b>

Calculation of  $(O - E)^2/E$

O	E	O - E	$(O - E)^2$	$(O - E)^2/E$
25	33.33	-8.33	69.39	2.082
35	33.33	1.67	2.79	0.083
40	33.33	6.67	44.49	1.335
40	33.33	6.67	44.49	1.335
24	33.33	-9.33	87.05	2.612
36	33.33	3.33	11.09	0.333
35	33.33	2.33	5.43	0.163
41	33.33	7.67	58.43	1.753
24	33.33	-9.33	87.05	2.612
			<b>Total</b>	$\chi^2 = 12.308$

Level of significance:  $\alpha = 0.05$

Degree of freedom:  $(r - 1)(c - 1) = (3 - 1)(3 - 1) = 4$

**MUQuestionPapers.com**

Critical Value: For 4 degree of freedom at 5% level of significance, the table value of  $x^2 = 9.488$

Decision: Since the calculated value  $x^2 = 12.308$  is more than the table value  $x^2 = 9.488$ , the hypothesis is not accepted.

There is a relation between the age and the choice of the vehicle.

**Q6 (a) State stokes theorem and evaluate  $\int [(x^2 + y^2)i + (x^2 - y^2)j]. d\vec{r}$  where C is the square in the xy – plane with vertices (1,0), (0,1), (-1,0) and (0,-1). (6M)**

**Solution:**

By Stokes theorem  $\int_c \vec{F} d\vec{r} = \iint_s \vec{N} \cdot \nabla \cdot \vec{F} ds$

$$\text{Now, } \nabla \times \vec{F} = \begin{vmatrix} i & j & k \\ \frac{\delta}{\delta x} & \frac{\delta}{\delta y} & \frac{\delta}{\delta z} \\ (x^2 + y^2) & (x^2 - y^2) & 0 \end{vmatrix} = (0 - 0)i - (0 - 0)j + (2x - 2y)k$$
$$= (2x - 2y)k$$

$$\vec{N} \cdot \nabla \cdot \vec{F} ds = (2x - 2y) dx \cdot dy$$

$$\iint_c \vec{N} \cdot \nabla \times \vec{F} \cdot ds = \iint_c (2x - 2y) \cdot dx \cdot dy \text{ where C is the square ABCD}$$

$$= 4 \iint_{\Delta OAB} (2x - 2y) \cdot dx \cdot dy$$

The equation of the line AB is  $\frac{y-1}{1-0} = \frac{x-0}{0-1}$  i.e.  $y = 1 - x$

$$\int_0^1 \int_{y=0}^{1-x} (2x - 2y) \cdot dy \cdot dx = \int_0^1 \left( 2xy - 2 * \frac{y^2}{2} \Big|_{y=0}^{1-x} \right) \cdot dx$$
$$= \int_0^1 2x(1-x) - 2 * \frac{(1-x)^2}{2} \cdot dx$$
$$= \int_0^1 2x - 2x^2 - 2 * \frac{(1-2x+x^2)}{2} \cdot dx$$
$$= \int_0^1 -3x^2 - 1 \cdot dx$$
$$= \left( -\frac{3x^3}{3} - x \Big|_{x=0}^1 \right)$$
$$= -2$$

**(b) Monthly salary X is an organisation is normally distributed with mean Rs. 3000 and standard deviation of Rs 250. What should be the normally minimum salary of an employee in this organisation so that the probability that an employee to top 5% employees? (6M)**

**Solution:**

$$\text{Mean}(m) = 3000$$

$$\text{Standard deviation } (\sigma) = 250$$

Let X denote monthly salary of a worker.

Let  $X_1$  be the minimum salary of the top 5% workers.  
 Let  $z_1$  be the corresponding pnv.

$$P(X > X_1) = 5$$

$$P(Z > Z_1) = 0.05$$

$$\begin{aligned} \therefore \text{Area between } z = 0 \text{ to } z = 5 \\ = 0.5 - 0.05 = 0.45 \end{aligned}$$

From table

$$z_1 = 1.645$$

$$\frac{X_1 - m}{\sigma} = 1.645$$

$$\frac{X_1 - 3000}{250} = 1.645$$

$$X_1 = 3411.25$$

Hence the minimum salary of the top 5% workers = Rs 3411.25

**(c) Using duality solve the following LPP**

**Maximize  $Z = 3x_1 + 2x_2$**

**Subject to  $2x_1 + x_2 \leq 5$**

$$x_1 + x_2 \leq 3$$

$$x_1, x_2 \geq 0.$$

**(8M)**

**Solution:**

The dual of the given problem is

Minimize  $w = 5y_1 + 3y_2$

Subject to  $2y_1 + y_2 \geq 5$

$$y_1 + y_2 \geq 3$$

$$y_1, y_2 \geq 0.$$

Introducing the slack and artificial variables, the problem becomes

Minimize  $w' = -w = -5y_1 - 3y_2$

$$\text{i.e. } w' = -5y_1 - 3y_2 - 0s_1 - 0s_2 - MA_1 - MA_2 \dots \dots \dots \text{(i)}$$

$$\text{subject to } 2y_1 + y_2 - s_1 - 0s_2 + A_1 - 0A_2 = 5 \dots \dots \dots \text{(ii)}$$

$$y_1 + y_2 - 0s_1 - s_2 - 0A_1 + A_2 = 3 \dots \dots \dots \text{(iii)}$$

Multiply (ii) and (iii) by M and add to (i)

$$w' = -5y_1 - 3y_2 + 3My_1 - 2My_2 - Ms_1 - Ms_2 - 0A_1 - 0A_2 - 5M$$

$$w' + (5 - 3M)y_1 + (3 - 2M)y_2 + Ms_1 + Ms_2 + 0A_1 + 0A_2 = -5M$$



Iteration Number	Basic Var.	Coefficient of						R.H.S. solution	Ratio
		$y_1$	$y_2$	$s_1$	$s_2$	$A_1$	$A_2$		
0	$w'$	$5-3M$	$3-2M$	$M$	$M$	0	0	$-5M$	
$A_1$ leaves	$A_1$	2*	1	-1	0	1	0	3	3/2
$y_1$ enter	$A_2$	1	1	0	-1	0	1	2	2
1	$w'$	0	$(1-M)/2$	$(5-M)/2$	$M$	$(3M-5)/2$	0	$-(15+M)/2$	
$A_2$ leaves	$y_1$	1	1/2	-1/2	0	1/2	0	3/2	3
$y_2$ enter	$A_2$	0	1/2*	1/2	-1	-1/2	1	1/2	1
2	$w'$	0	0	2	1	$M-2$	$M-1$	-8	
	$y_1$	1	0	-1	1	1	-1	1	
	$y_2$	0	1	1	-2	-1	1	1	

Since,  $s_1 = 2$ ,  $s_2 = 1$  and  $w'_{\max} = -8$ , therefore  $w'_{\min} = 8$

$x_1 = 2$ ,  $x_2 = 1$  and  $z_{\max} = -8$ .