

(3hours)

[Total marks: 80]

- N.B.** (1) Question No. 1 is compulsory.
 (2) Answer **any Three** from remaining
 (3) Figures to the right indicate full marks.

1. (a) Find Laplace transform of $e^{-4t} \sin ht \sin t$. 5
- (b) Does there exist an analytic function whose real part is $x^3 - 3x^2y - y^3$. Give justification. 5
- (c) Show that $\{\cos x, \cos 2x, \cos 3x, \dots\}$ is a set of orthogonal functions over an interval $(-\pi, \pi)$. 5
- (d) Evaluate $\int_0^{2+i} z^2 dz$ along the line joining the point $z_1 = 0$ and $z_2 = 2 + i$. 5
2. (a) Obtain the Taylor's and Laurent series which represent the function,
 $f(z) = \frac{1}{(z+1)(z+3)}$ valid in the regions,
 (i) $|z| < 1$ (ii) $1 < |z| < 3$ (iii) $|z| > 3$ 6
- (b) Find the bilinear transformation which maps the points $z = \infty, i, 0$ into the points $w = 0, i, \infty$. 6
- (c) Using Laplace transform, solve the differential equation :
 $\frac{d^2x}{dt^2} + 4x = t$ with $x(0) = 1, x'(0) = -2$ 8
3. (a) Solve $\frac{\partial^2 u}{\partial x^2} - 2 \frac{\partial u}{\partial t} = 0$ by Bender-Schmidt method, given
 $u(0, t) = 0, u(x, 0) = x(4 - x), u(4, t) = 0$, assuming $h = 1$, find u upto $t=5$. 6
- (b) Using convolution theorem find the inverse Laplace transform of
 $\frac{s}{(s^2 + 1)(s^2 + 4)}$. 6
- (c) Determine the solution of one-dimensional heat equation $\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$
 under boundary condition $u(0, t) = u(l, t) = 0, u(x, 0) = x$, l being the length of rod. 8

[TURN OVER]

4. (a) Using Residue theorem, evaluate, $\int_0^{2\pi} \frac{d\theta}{5 + 3\sin \theta}$. 6

(b) Find the inverse Laplace transform of the following:

$$\frac{s^2 + 2s + 3}{(s^2 + 2s + 2)(s^2 + 2s + 5)}$$
 6

(c) Obtain Half Range Sine Series of $f(x) = x(\pi - x)$ in $(0, \pi)$.

Hence, evaluate $-\sum_{m=0}^{\infty} \frac{(-1)^m}{(2m+1)^3}$.

8

5. (a) If $f(x) = e^{-3x}$, $-1 < x < 1$. Obtain Complex form of $f(x)$ in $(-1, 1)$. 6

(b) Find the orthogonal trajectory of the family of curves $3x^2y - y^3 = c$. 6

(c) Solve by Crank-Nicholson simplified formula $\frac{\partial^2 u}{\partial x^2} - \frac{\partial u}{\partial t} = 0$,

$$u(0, t) = 0, u(1, t) = 2t, u = 0, \text{ for two time steps taking } h = 0.25. \quad 8$$

$$u(x, 0) = 0$$

6. (a) Obtain the Fourier series for $f(x)$ where

$$f(x) = x + \frac{\pi}{2} \quad -\pi < x < 0$$

$$= \frac{\pi}{2} - x \quad 0 < x < \pi$$

(b) Prove that $\int_0^{\infty} e^{-t} \frac{\sin^2 t}{t} dt = \frac{1}{4} \log 5$ 6

(c) Find bilinear transformation which maps the points $z = 1, i, -1$ onto the points $w = i, 0, -1$. Hence, find the image of $|z| \leq 1$ onto the w -plane. 8