(3 Hours) (Total Marks: 100)

## Please check whether you have the right question paper.

.Б.:	2) Figures to the Right indicate full marks	3					
	3) Use of log table/non-programmable calculator is allowed.	7,26					
Q.1.	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~						
A)	Distinguish between:						
	a) polarizable electrode and non-polarizable electrode	ST.					
	b) voltammetry and polarography						
<b>B</b> )	Explain the need to remove dissolved oxygen from a solution before recording its						
	polarogram.						
<b>C</b> )	A Zn <sup>+2</sup> ion solution of unknown concentration gave a diffusion current of 4.25 μA. When	4					
	3.0 cm <sup>3</sup> of 0.01 mol dm <sup>-3</sup> solution of ion was added to 25.0 cm <sup>3</sup> of unknown, the diffusion						
	current increased to $12.5\mu A$ . Calculate the concentration of $Zn^{+2}ions$ in the sample						
	solution.						
D)	A 0.878 mM Pb <sup>+2</sup> ion solution gave a diffusion current of 7.75 μA. The mercury flow rate						
	and drop lifetime were 2.63 mgs <sup>-1</sup> and 2.88 s respectively. What is the diffusion						
	coefficient of Pb <sup>+2</sup> ?						
E)	Give the advantages and limitations of amperometric titrations.						
F)	Explain: a) basic principle of amperometric titrations.	4					
	b) draw a neat labelled diagram of the H-shaped polarographic cell.						
Q.2.	Attempt any four of the following:						
A) (	Mention the detectors used in GC. Describe the principle and construction of any one						
	detector with a neat labelled diagram.						
<b>B</b> )	What are the applications of gas chromatography?	4					
<b>C</b> )	Explain the terms w.r.t. GC: retention time, HETP and resolution.						
<b>D</b> )	Substances A and B gave retention time 15.15 and 16.60 min. respectively, on a 0.2m	4					
300	column. An unretained species passed through the column in 1.30 min. The peak widths						
	at base for A and B were 1.01 and 1.20 min. respectively. Calculate i) resolution						
	ii) number of theoretical plates iii) plate height.						
<b>E</b> )	What is ion exchange capacity? How is cation exchange capacity of a resin						
	experimentally determined?						
<b>F</b> )	Define selectivity coefficient for an ion exchange process. What are the factors affecting	4					
S S S S	selectivity coefficient?						
	XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX						

66348 Page 1 of 4

Q.3	Attempt any four of the following:	ئے کر اگر				
A	Discuss the use of chemicals as food preservatives.	5				
В	What is the role of chicory in coffee? Discuss the constituents present in coffee.					
C	Write the nutrients present in milk and discuss the analysis of milk for lactose by Lane	5				
	Eynon's method.	86				
D	Write the composition of tea and discuss the detection of any four adulterants present in it.	5				
$\mathbf{E}$	Write the properties of antiperspirants and describe a gravimetric method to estimate	5				
	amount of zinc in antiperspirants.	257				
F	What are sensory properties of cosmetics? Describe a method to estimate amount of	5				
	magnesium present in face powder.					
<b>Q.4</b>	Attempt any four of the following:					
A	Draw a schematic diagram of TGA instrument and discuss any three of its components.	5				
В	Discuss the principle of DTA and explain the nature of DTA curve for dehydration of	5				
	CuSO <sub>4</sub> . 5H <sub>2</sub> O.					
C	Discuss the factors that influence the TG curve.	5				
D	a) Compare the technique of TGA with DTA.	3				
	b) Write a note on: Reference material used DTA.	2				
E	Discuss the principle of thermometric titrations and explain its application in the titration	5				
	of HCl against NaOH.					
F	Discuss the need for validation of a method and explain the significance of linearity,	5				
	selectivity and specificity w.r.t. method validation.					
Q.5.A	State true or false (any five)	5				
a)	The drop time can be changed by adjusting the height of the mercury reservoir in					
25 TO V	dropping mercury electrode.					
<b>b</b> )	In amperometric titrations, a 'V' shaped titration curve is obtained, when only the titrant					
	is reducible species.					
(c)	Limiting current is the sum of diffusion current and residual current.					
<b>d</b> )	Rotating platinum electrode is used for determination of halides by precipitation titration.					
e)	The indifferent electrolyte present in large concentration in the polarographic analysis is					
	known as the maxima suppressor.					
<b>f</b> )	Polarogram is a plot of current versus voltage.					
<b>g</b> )	Due to continuous renewal of DME, simultaneous determination of several metals is					
	possible.					
<b>h</b> )	The rotating platinum electrode has the working range upto +0.4V versus SCE.					

Page 2 of 4

Q.5.B	Select the correct option: (any five)	75 75
a)	The contribution to band broadening due to different velocities of solute molecules across	35
	the band is known as	30
	i)Eddy diffusion ii) Longitudinal diffusion iii) Non equilibrium mass transfer	
<b>b</b> )	For weakly basic anion exchangers, pH must be	25
	i)greater than 7 ii) less than 7 iii) 7	ST.
<b>c</b> )	Cation exchange resin is a polymer containing groups as an integral part of the	
	resin.	
	i)amino ii) sulphonic iii) quarternary amino	
d)	Sample port temperature in GC is kept at the boiling point of the	
	volatile component.	
	i)most ii) least iii) neither (i) nor (ii)	
e)	Ion exchange chromatography can be used for	
	i) demineralisation of water ii) separation of amino acid iii) both (i) and (ii)	
f)	Swelling of a resin is due to high proportion of groups	
	i)polar ii) non-polar iii) neutral	
g)	is a natural ion-exchanger.	
	i) clay ii) Amberlite IR-120 iii) Dowex-3	
Q.5C	State true or false (any five)	5
a) (1)	Irradiation reduces the need of food preservatives.	
<b>b</b> )	Mannitol is used as an indicator for the determination of boric acid in food sample	
50 00 00 00 00 00 00 00 00 00 00 00 00 0	titrimetrically.	
<b>c</b> )	The action of sodium salt of benzoic acid as preservative is pH dependent.	
<b>d</b> )	Castrol oil is used in lipstick for the dissolution of dyes.	
<b>e</b> )	Zinc oxide is added to face powder to give a covering power.	
<b>f</b> )	Aluminium salts are used in antiperspirants to increase its fragrance.	
<b>g</b> )	Sugar syrup is the adulterant added to honey.	
h	Amount of kaolin in face powder should not exceed 25%.	

66348 Page 3 of 4

<b>O.5D</b>	Select the	correct	correct	ontion	(onv	fivo)	
U.SD	Select the	correct	correct	opuon:	(anv	nve	ı

a) In the application of DTA, which of the following parameter is measured for glasses?i)concentration ii)cooling temperature iii)transition temperature

5

b) Which of the following thermal technique detects every physical or chemical change whether or not accompanied by change in weight?

i)TGA ii)DTA iii)both DTA and TGA

- c) Which of the following represents the loss of water of crystallisation in the TGA curve?

  i)downward slope ii)upward slope iii)upward peak.
- d) How many peaks are observed in the thermometric titration curve for the titration of mixture Ca<sup>+2</sup> and Mg<sup>+2</sup> against EDTA?

i)one ii)two iii)three

- e) The formation of which of the following product shows an upward peak in the DTA curve of calcium oxalate monohydrate in the atmosphere of air?

  i)calcium oxalate ii)calcium carbonate iii)calcium oxide iv) calcium
- f) How many endotherms are observed in the DTA curve of calcium oxalate monohydrate?

  i)one ii)two iii)three iv)four
- g) Which of the following term is used for expressing stability of a method against extraneous influencing parameters.

i) precision ii) ruggedness iii)accuracy iv) selectivity

66348 Page 4 of 4