

Duration:[3 Hours]

[Total Marks: 100]

- N.B. 1) All questions are compulsory.
2) Figures to the right indicate full marks.

1. Choose correct alternative in each of the following: (20)

- i. Let c be a proper coloring of $G = (V, E)$ using t colors, then the coloring partitions V into
 - (a) $t - 1$ parts
 - (b) t parts
 - (c) one part
 - (d) 2 parts
- ii. If graph G is k -critical then
 - (a) G is acyclic
 - (b) G is disconnected
 - (c) G is connected
 - (d) all of these.
- iii. Which of the following can be a chromatic polynomial?
 - (a) $k^4 - 3k^3 + 3k^2 - k$
 - (b) $3k^3 - 4k^2 + k$
 - (c) $k^4 - 5k^3 + 7k^2 - 6k + 3$
 - (d) $k^3 + k^2 + k$
- iv. K_n is planar if
 - (a) $n > 4$
 - (b) $n \leq 4$
 - (c) $n = 5$
 - (d) None of these
- v. A connected planar graph has an equal number of vertices and faces. If there are 20 edges in this graph, the number of vertices must be:
 - (a) 9
 - (b) 10
 - (c) 20
 - (d) 11
- vi. If f is a flow in a network N and P be any f -augmenting path with tolerance $\epsilon(P) > 0$, then define a new flow f' as follows : $f'(a) = f(a) + \epsilon(P)$ for an forward arc $a \in P$, $f'(a) = f(a)$ for an backward arc $a \in P$ and $f(a) = f(a)$ for other arcs a of N . Then value of f' equals to
 - (a) $val f + \epsilon(P)$
 - (b) $val f - \epsilon(P)$
 - (c) same as $val f$
 - (d) $val f \times \epsilon(P)$
- vii. Let $R(x, B)$ denotes the rook polynomial for the board B of darkened squares consisting of m rows and n columns, then
 - (a) constant term is 1
 - (b) coefficient of x^k is number of ways of placing k non capturing rooks
 - (c) $r_k(B) = 0$ if $k > \min\{m, n\}$
 - (d) all of the above.
- viii. The number of ways to climb a staircase with 12 steps taking 1 or 2 steps at a time is
 - (a) 987
 - (b) 610
 - (c) 377
 - (d) 233

[TURN OVER

- ix. The number of different system of distinct representatives for the family $A_1 = \{2, 3, 4\}$, $A_2 = \{1, 3, 4\}$, $A_3 = \{1, 2, 4\}$, $A_4 = \{1, 2, 3\}$ is
 (a) 4! (b) 3! (c) 9 (d) None of these
- x. If M is maximum matching then which one of the following statement is true?
 (a) There does not exists any matching M' such that $|M'| < |M|$
 (b) There does not exists any matching M' such that $|M'| > |M|$
 (c) There exists any matching M' such that $|M'| > |M|$
 (d) None of these

2. (a) Attempt any **ONE** question from the following: (8)

- i. For a simple graph G of order p and size q , prove that $\pi_k(G)$, the chromatic polynomial of the graph G , is monic polynomial of degree p in k with integer coefficients and constant term zero. Further prove that its coefficients are alternate in sign and the coefficient of k^{p-1} is $-q$.
- ii. For any simple graph G , prove that $\kappa(G) \leq \kappa'(G) \leq \delta(G)$ where $\kappa(G)$ denote the vertex connectivity and $\kappa'(G)$ denotes the edge connectivity and $\delta(G)$ denotes the minimum degree of a graph G .

(b) Attempt any **TWO** questions from the following: (12)

- i. State Vizing theorem for edge coloring of graphs. Show that $\chi'(G) \geq \Delta(G)$ where $\chi'(G)$ denotes edge chromatic number and $\Delta(G)$ denotes the maximum degree of G . Give an example of the graph for which $\chi'(G) = \Delta(G)$.
- ii. Let $\pi_k(G)$ denote the chromatic polynomial of the graph G . If G is simple graph then prove that $\pi_k(G) = \pi_k(G - e) - \pi_k(G.e)$ where e is an edge of G .
- iii. If G is a graph with p vertices and \overline{G} is its complement of G , then show that $\chi(G) + \chi(\overline{G}) \leq p + 1$, where $\chi(G)$ is the vertex chromatic number of graph G .
- iv. If G is a (p, q) graph, then prove that $\chi(G) \geq \frac{p^2}{p^2 - 2q}$ where $\chi(G)$ denotes the vertex chromatic number of G .

3. (a) Attempt any **ONE** question from the following: (8)

- i. Show that every planar graph is 5 vertex colorable.
- ii. State and prove Max Flow - Min Cut Theorem.

(b) Attempt any **TWO** questions from the following: (12)

- i. Show that there is at least one face of every polyhedron is bounded by an n -cycle for some $n = 3, 4$ or 5 .
- ii. Show that the edge e is a loop in G if and only if e^* is a bridge in G^* where G^* is dual of graph G .

[TURN OVER

- iii. Define a value of flow and capacity of cut in network N . If f is any flow and K be any cut in a network N then show that $val(f) \leq cap(K)$.
- iv. If G is a connected simple planar graph with $p \geq 3$ vertices, q edges and f regions then
 - I) Show that if $q = 3p - 6$ then each region is triangle.
 - II) Deduce that a convex polyhedron with 12 vertices and 20 faces is composed entirely of triangles.

4. (a) Attempt any **ONE** question from the following: (8)

- i. State and prove the necessary and sufficient condition for a family of n sets to have System of Distinct Representative.
- ii. An elf has a staircase of n stairs to climb. Each step it takes can cover either one stair or two stairs. Find a recurrence relation for a_n , the number of different ways for the elf to ascend the n -stair staircase and solve it by using generating function.

(b) Attempt any **TWO** questions from the following: (12)

- i. Define a rook polynomial. Let $R_{n,m}(x)$ be the rook polynomial for the $n \times m$ chess board, all squares may have rooks. Show that $R_{n,m}(x) = R_{n-1,m}(x) + mxR_{n-1,m-1}(x)$
- ii. Show that a matching M in G is a maximum matching if and only if G contains no M -augmenting path.
- iii. Find the coefficient of x^{16} in $(x^2 + x^3 + x^4 + \dots)^5$. What is the coefficient of x^r ?
- iv. Let h_n denote the number of nonnegative integral solutions of the equation $3e_1 + 4e_2 + 2e_3 + 5e_4 = n$. Find the generating function $f(x)$ for $h_0, h_1, \dots, h_n, \dots$

5. Attempt any **FOUR** questions from the following: (20)

- (a) For any graph G , prove that $\chi(G) \leq \Delta(G) + 1$ where $\chi(G)$ represents vertex chromatic number of a graph G and $\Delta(G)$ denotes the maximum degree of G . Give an example of graphs for which $\chi(G) < \Delta(G)$.
- (b) Prove that every tree with $n \geq 2$ vertices is 2-chromatic.
- (c) If G be a simple connected graph with at least 11 vertices then prove that either G or its complement \bar{G} must be nonplanar.
- (d) If f is flow in a network N and P is any f -incrementing path, then show that there exists a revised flow f' such that $val f' > val f$.
- (e) Find the rook polynomial for the following $\{(1, 1), (2, 5), (3, 3), (4, 2), (4, 4), (5, 1), (5, 3)\}$.
- (f) Let $\{A_1, A_2, \dots, A_n\}$ be a family of sets such that for each $k, 1 \leq k \leq n$ and for each choice of $1 \leq i_1 < i_2 < \dots < i_k \leq n, |A_{i_1} \cup A_{i_2} \cup \dots \cup A_{i_k}| \geq k + 1$. Let x be any element of A_1 . Show that $\{A_1, A_2, \dots, A_n\}$ has a system of distinct representatives in which x represents A_1 .
